

CONCLUSION

This study analyzed the effects of a new combined technique for treating adipose tissue disorders such as gynoid lipodystrophy and localized fat. The results have shown that the technique is safe, effective, and well tolerated in non-invasive bodily procedures. In particular, the authors believe that the ideal patients for this type of procedure are those with localized fat and cellulite, rather than those seeking to lose weight. This technique would also be an option for treating patients

who are advised not to have surgical treatment. Laboratory analysis showed that no significant alterations of serum fats, blood glucose levels, or liver function are caused by this procedure. It is always worth highlighting that a comprehensive patient history that describes previous diseases, accompanied by a detailed physical examination and an analysis of the patient's level of expectations, are vital for success and to avoid patient dissatisfaction. ●

REFERENCES

1. Goldman MP. Cellulite a review of current treatments. *Cosmet Dermatol.* 2002; 15:17-20.
2. Lucas MRA. Association of multipolar radiofrequency with vacuum massage and pulsed cavitation ultrasound. *South Am J Aesthet Med.* 2011;07:08-11.
3. Palumbo P, Cinque B, Miconi G, La Torre C, Zoccali G, Vrentzos N, et al. Biological effects of low frequency high intensity ultrasound application on ex vivo human adipose tissue. *Int J Immunopathol Pharmacol.* 2011;24(2):411-22.
4. Boisnic S, Branchet MC. Ex vivo human skin evaluation of localized fat reduction and anti-aging effect by TriPollar radio frequency treatments. *J Cosmet Laser Ther.* 2010;12(1):25-31.
5. Sousa AP, Santos JN, Reis JAJR, Ramos TA, Souza J, Cangussú MC, et al. Effect of LED phototherapy of three distinct wavelengths on fibroblasts on wound healing: a histological study in a rodent model. *Photomed Laser Surg.* 2010;28(4):547-52.
6. Chang P, Wiseman J, Jacoby T, Salisbury AV, Ersek RA. Noninvasive mechanical body contouring (Endermologie): a one-year clinical outcome study update. *Aesthetic Plast Surg.* 1998; 22(2): 145-53.
7. Huang PJ, Huang YC, Su MF, Yang TY, Huang TY, Huang JR, et al. *In vitro* observations on the influence of copper peptide aids for the LED photoirradiation of fibroblast collagen synthesis. *Photomed Laser Surg.* 2007;25(3):183-90.
8. van Hees AM, Jocken JW, Essers Y, Roche HM, Saris WH, Blaak EE. Adipose triglyceride lipase and hormone-sensitive lipase protein expression in subcutaneous adipose tissue is decreased after an isoenergetic low-fat high-complex carbohydrate diet in the metabolic syndrome. *Metabolism.* 2012 Apr 30. [Epub ahead of print].
9. Enjoji M, Yasutake K, Kohjima M, Nakamuta M. Nutrition and nonalcoholic Fatty liver disease: the significance of cholesterol. *Int J Hepatol.* 2012;2012:925807.
10. Weber K, Erben RG. Differences in triglyceride and cholesterol metabolism and resistance to obesity in male and female vitamin D receptor knockout mice. *J Anim Physiol Anim Nutr (Berl).* 2012 Apr 30. [Epub ahead of print].
11. Scandurra S. Analysis and overview about the innovative method the "stable cavitation" for treatment of localized adiposity and oedema-fibrosclerotic panniculopathy. Use of generator at "cavitation" dominance. Monitoring and follow-up of blood and urine parameters on 3 groups of 7 patients each. *Plast Aesthetic Surg.* 2005: 1-9.
12. Neira R, Arroyave J, Ramirez H, Ortiz CL, Solarte E, Sequeda F, et al. Fat liquefaction: Effect of low-level laser energy on adipose tissue. *Plast Reconstr Surg.* 2002; 110(3): 912-22.
13. Zhou Y, Yang K, Cui J, Ye JY, Deng CX. Controlled permeation of cell membrane by single bubble acoustic cavitation. *J Control Release.* 2012; 157(1): 103-11.
14. Royo de la Torre J, Moreno-Moraga J, Muñoz E, Cornejo Navarro P. Multisource, Phase-controlled Radiofrequency for Treatment of Skin Laxity: Correlation Between Clinical and In-vivo Confocal Microscopy Results and Real-Time Thermal Changes. *J Clin Aesthet Dermatol.* 2011;4(1):28-35.
15. Mlosek RK, Woźniak W, Malinowska S, Lewandowski M, Nowicki A. The effectiveness of anticellulite treatment using tripolar radiofrequency monitored by classic and high-frequency ultrasound. *J Eur Acad Dermatol Venereol.* 2012;26(6):696-703.
16. Lee SY, You CE, Park MY. Blue and red light combination LED phototherapy for acne vulgaris in patients with skin phototype IV. *Lasers Surg Med.* 2007;39(2):180-8.
17. Weiss RA, McDaniel DH, Geronemus RG, Weiss MA, Beasley KL, Munavalli GM, et al. Clinical experience with light-emitting diode (LED) photomodulation. *Dermatol Surg.* 2005;31(9 Pt 2):1199-205.
18. Shack RB. Endermologie: taking a closer look. *Aesthet Surg J.* 2001;21(3):259-60.
19. Latrenta GS, Mick SL. Endermologie after external ultrasound-assisted lipoplasty (EUAL) versus EUAL alone. *Aesthet Surg J.* 2001;21(2):128-35.